

Errata: Corrections to “Review of the genus *Carinapex* Dall, 1924 with the description of ten new species (Gastropoda: Conoidea: Horaiclavidae) from the Pacific Ocean”

Shawn G. Wiedrick

An article by Shawn G. Wiedrick in Volume 47(1) of the *Festivus* included a review of the genus *Carinapex* Dall, 1924 with descriptions of new taxa from the Indo-Pacific. Two similar *Carinapex* species were compared, *C. minutissima* Garrett, 1873 (fig. 1) and *C. amirowlandae* Wiedrick, 2015 (fig. 3) with the distribution map (fig. 2) of *C. minutissima* inadvertently omitted from the article. It is important to note the similarities of the two species which can easily be confused based on the overlap of distribution within both ranges and their morphological affinities. After reviewing 118 lots from the Natural History Museum of Los Angeles County and the Shawn Wiedrick collection, which included over 3080 specimens, it was concluded that two separate species exist. Upon close morphological inspection, *C. minutissima* is primarily much smaller and has one to one-and-a-half initial protoconch whorls, white in color. *C. amirowlandae* is relatively larger, with first two protoconch whorls brown, both species easily distinguishable in fresh specimens. Equally as important is the distribution of *C. minutissima*, which expands to extreme localities (Red Sea and Hawai‘i) which are completely absent in the range of *C. amirowlandae* (fig. 4). Additionally, plate 2, figure 39-41, 46 should read *Carinapex mooreorum*, not *C. moorei*.



Figure 1. *Carinapex minutissima* (Garrett, 1873), Hawai‘i, USA, height 3.1 mm, width 1.2 mm.



Figure 3. *Carinapex amirowlandae* Wiedrick, 2015, Mactan Island, Philippines, height 3.1 mm, width 1.2 mm.




Figure 2. Distribution of *Carinapex minutissima* Garrett, 1873

- Exclusively *Carinapex minutissima* in relation to *C. amirowlandae* n. sp.
- 50-99% *Carinapex minutissima* in relation to *C. amirowlandae* n. sp.
- 25-49% *Carinapex minutissima* in relation to *C. amirowlandae* n. sp.
- 1-24% *Carinapex minutissima* in relation to *C. amirowlandae* n. sp.




Figure 4. Distribution of *C. amirowlandae* Wiedrick, 2015

- Exclusively *Carinapex amirowlandae* in relation to *C. minutissima* n. sp.
- 50-99% *Carinapex amirowlandae* in relation to *C. minutissima* n. sp.
- 25-49% *Carinapex amirowlandae* in relation to *C. minutissima* n. sp.
- 1-24% *Carinapex amirowlandae* in relation to *C. minutissima* n. sp.